

Save on QUVIVIQ

A prescription medication to treat adults who have trouble falling asleep or staying asleep (insomnia).

You pick up the prescription, QUVIVIQ will help take care of the rest.

Simply cut out and bring this copay card with you when picking up your first prescription.

RxBIN: 600426 RxGRP: EC44101001 ID: 09776180081 RxPCN: 54







Get your first 30-day supply of QUVIVIQ for as little as \$0

and refills for as little as \$25*

*For eligible commercially insured patients. Terms and conditions apply. Please read the full terms and conditions.

CUT ALONG THE DOTTED LINE -

CUT ALONG THE DOTTED LINE

Restrictions: This offer is not valid for Cash-Paying Patinswed by commercial insurance and your prescription insurance overland to an oto event full cost of your prescription (2) you do not have prescription insurance coverages through a state of federal healthcare programs, including but not intended to Medicare Part D. Medicaid, Medigap, Veterana Affairs (VA), or Department of Defense (DOD) programs; patients who move from commercial insurance plans to state or federal healthcare programs will not longer be eligible; (3) you are 18 years of age or older, (4) you are a resident of the United States or Puerto Rico.

Medicare Part D, Medicaid, Medigap, Veterans Affairs (VA), or begrattement of Defense (DOD) programs; patients who move from commercial insurance plans to state or federal healthcare organs will no longer be eligible; (3) you are 18 years of age or older, (4) you are a resident of the United States or Puerto Rico.

TERMS OF USE: Eligible commercially insured patients with a valid prescription for QUVIVIQ (daridorexant) 25 mg and 50 mg hop present this savings card at participating pharmacies may pay as little as 90 for their first fill, and 925 for subsequent fills or 30-day supply. Eligible insurance of 100 commercially insured patients with a spiles; patient out-of-pocket expenses may vary. The patient is responsible for applicable taxes—void where prohibited by law, and, or restricted, and other restrictions may apply. Patients, pharmacists, and prescribers cannot seek reimbursement from the prohibited of the promote or any third party for any part of the benefit reserved to the promote or any third party for any part of the benefit reserved to the promote or amend this offer, eligibility, and terms of use at any time without notice. Must present offer along with a valid prescription at the time of purchased, or some control of the promote of





Join Q&You today

Get more info on savings, support on taking QUVIVIQ (daridorexant), sleep tips, and more with Q&You at QUVIVIQ.com/join.

If you have any questions, contact us at 877-478-8484 or by live chatting at vitacarerx.com. (Hours of Operation: M-F 8AM-12AM ET, S & S 9AM-5PM ET)

Indication

QUVIVIQ is a prescription medicine for adults who have trouble falling asleep or staying asleep (insomnia).

Important Safety Information

Do not take QUVIVIQ if you fall asleep often at unexpected times (narcolepsy).

Please see additional Important Safety Information on the following page and full Prescribing Information and Medication Guide.

Important Safety Information (cont'd)

QUVIVIQ may cause serious side effects, including:

- Decreased awareness and alertness.
 The morning after you take QUVIVIQ, your ability to drive safely and think clearly may be decreased. You may also have sleepiness during the day.
 - Do not take more QUVIVIQ than prescribed.
 - Do not take QUVIVIQ unless you are able to stay in bed for at least 7 hours before you must be active again.
 - Take QUVIVIQ at night within 30 minutes before going to bed.

QUVIVIQ is a federally controlled substance because it can be abused or lead to dependence.

Before taking QUVIVIQ, tell your healthcare provider about all of your medical conditions, including if you:

- have a history of depression, mental illness, or suicidal thoughts or actions; drug or alcohol abuse or addiction; a sudden onset of muscle weakness (cataplexy); daytime sleepiness
- have lung or breathing problems, including sleep apnea
- have liver problems
- are pregnant or plan to become pregnant
- are breastfeeding or plan to breastfeed

Tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

- Taking QUVIVIQ with certain medicines can cause serious side effects. QUVIVIQ may affect the way other medicines work and other medicines may affect the way QUVIVIQ works.
- Do not take QUVIVIQ with other medicines that can make you sleepy unless instructed by your healthcare provider.

Please see full <u>Prescribing Information</u> and Medication Guide.

What should I avoid while taking QUVIVIQ?

- Do not drink alcohol while taking QUVIVIQ.
 It can increase the effects of alcohol, which can be dangerous.
- Do not drive, operate heavy machinery, do anything dangerous, or do other activities that require clear thinking if you do not feel fully awake, or you have taken QUVIVIQ and have less than a full night of sleep (at least 7 hours), or if you have taken more QUVIVIQ than prescribed.

QUVIVIQ may cause other serious side effects, including:

- Worsening depression and suicidal thoughts. Call your healthcare provider right away if you have any worsening depression or thoughts of suicide or dying.
- Temporary inability to move or talk (sleep paralysis) for up to several minutes, or hallucinations while you are going to sleep or waking up.
- Complex sleep behaviors such as sleepwalking, sleep-driving, preparing and eating food, making phone calls, having sex or doing other activities while not fully awake that you may not remember the next morning. Stop taking QUVIVIQ and call your healthcare provider right away if you experience a complex sleep behavior.

The most common side effects of QUVIVIQ are headache and sleepiness.

These are not the only side effects of QUVIVIQ. Call your doctor for advice about side effects.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.



